

FATHERS INVOLVEMENT AND PARTICIPATION IN FEEDING THE UNDER FIVES;

CAN IT SAVE THE CHILDRENS NUTRITIONAL STATUS?

Kaggwa Margaret

Uganda

Fathers should be involved in feeding to improve on the nutrition status of the under fives. Are the bread winners in most families hence the need to involve them in nutrition of fewer than fives. Nutritional status is determined by three broad factors i.e. food, care and health. Since fathers are the sole provider for family, lack of resources by the care taker to cater for health will affect nutrition of under fives. Malnutrition if not, prevented, treated in early stages promptly, and rehabilitee becomes permanent cause brain damage in the under-fives. Father involved in child neglect, abandonment, divorce, inequalities are leading secondly malnutrition in low income countries. Low adequate feeding practices i.e. continued breast feeding, adequate feeding frequency and diversity in children 6-23 months: 37 percent *ZDHS 2007*). Lack of resources or knowledge of care takers, along with illnesses such as diarrhea, pneumonia, malaria, and HIV and AIDS, often exacerbated by intestinal parasites, are immediate causes of malnutrition. Underlying and more basic causes include poverty, household food insecurity, unsanitary health environment, illiteracy, social norms, and emergencies In England, the Government's strategy for securing parental involvement was first set out in the 1997 White Paper, Parent & under-fives. Therefore fathers are

CAN IT SAVE THE CHILDRENS NUTRITIONAL STATUS ?

Kaggwa Margaret

Uganda

The silent component neglected," men involvement and participation", untapped targeting only women of child bearing age yet men the policy maker, researchers, bread earner ,administrator, financier, leader ,and God given gift and control all the environment. Malnutrition if not, prevented, treated in early stages promptly, and rehabilitee becomes permanent cause brain damage in the under-fives. Men involved in child neglect, abandonment, divorce, inequalities are leading secondly malnutrition in low income countries. Low adequate feeding practices i.e. continued breast feeding, adequate feeding frequency and diversity in children 6-23 months: 37 percent *ZDHS 2007*). Yet under nutrition is determined by a number of factors which operate at different levels. An individual's nutritional status is influenced by three broad categories of factors -- food, care, and health -- and adequate nutrition requires the presence of all three. Poor infant and young child feeding practices due to lack of resources or knowledge of care takers, along with illnesses such as diarrhea, pneumonia, malaria, and HIV and AIDS, often exacerbated by intestinal parasites, are immediate causes of malnutrition. Underlying and more basic causes include poverty, household food insecurity, unsanitary health environment, illiteracy, social norms, and emergencies In England, the

Government's strategy for securing parental involvement was first set out in the 1997 White Paper, Parent & under-fives. Therefore men should be key in reduction to the nutritional status of the under-fives at all levels.

Globally 161 million are under-five have child malnutrition .11 million in Africa and 4 million in Latin America and the Caribbean. High prevalence levels of stunting among children under-five years of age in Africa (United States of America), Ethiopia is one of the countries with the highest levels of malnutrition particularly prevalent among children under five years of age (UNICEF) 2006. The World Health Organization predicts that type 2 diabetes will become the seventh largest killer by 2030. Globally, 1.4 billion people are overweight and obese, while 870 million suffer from chronic hunger, representing a dual burden of malnutrition. Australia is by no means the 'fattest' country. Twenty-five per cent of children in Australia were deemed overweight and obese in 2011-12.

Prevalence of vitamin A deficiency: 54 percent in children under five and 13 percent among women of child bearing age;

Prevalence of anemia: 53 percent in children under five and 30 percent in women of child bearing age (*Source: NFNC2002*)

Under-nutrition can be greatly reduced through the delivery of simple interventions at key stages of the life cycle - for the mother: before she becomes pregnant, during pregnancy, and while breastfeeding; for the child, in infancy and early childhood. Maternal under-nutrition leads to intra-uterine growth retardation and low birth weight. Malnourished children have substantially lower chances of survival than children who are well nourished. They are much more likely to suffer from a serious infection and die from common childhood diseases such as diarrhoea, pneumonia, measles, and malaria. Maternal and child under-nutrition is estimated to contribute to one-third of child mortality.

A package of effective nutrition interventions to reduce the levels of chronic malnutrition (stunting) has been globally agreed upon by experts in the field. It includes: adequate maternal nutrition during pregnancy and lactation, early initiation of breastfeeding, exclusive breast feeding for the first 6 months, continued breastfeeding and adequate complementary feeding from 6 to 24 months, and increased micronutrient intakes during the critical 1,000 days.

UNICEF/World Bank/WHO/GAVI/MoH supports the delivery of priority interventions at different stages of the life cycle: policy development, system strengthening, service delivery, advocacy, capability building, partnerships, resource mobilization, innovations, and knowledge management

Provides technical and financial support to the Ministry of Health under the following programs:
- Bi-annual Child Health Weeks for vitamin A supplementation and de-worming. Children under 5 receive vitamin A capsules and de-worming tablets.

Supports planning, supervision, and implementation.
- Promotion of adequate infant and young child feeding practices through the development of appropriate guidelines and training materials, and capacity building of health workers at facility

and community level.

- Management of severe acute malnutrition through the provision of therapeutic nutrition foods, medicines, anthropometric equipment, as well as training of health care workers.

Nutrition is under the program me result, “Children, mothers, and pregnant women benefit from high impact interventions contributing to the attainment of Millennium Development Goal targets for child and maternal survival and development.”

Primary Health Care,Almata Decra lation 1987 targeting by the year 2000 passed unnoticed little was done most communities.

A purposive study was done among men using the same questionnaire in Katanga peri urban slum 2003 (30%), Mulago medical workers 2007 (27%) and at Komamboga rural health center 2016 (50%) about the involvement of males and their participation in feeding the under-fives revealed out that both elite , slum dwellers and rural health workers had never asked the number of times mothers breast feed the baby, all admitted to buy foods in house but didn’t know the weaning foods, they admitted to enjoy the fun, love their babies and return home late when babies are asleep rarely earlier than they sleep. At the nutritional Unit of Mulago Kampala national referral hospital out of the 20 admitted cases per day only one mother is being escorted by the husband. Relapses of malnutrition have been experienced by the unit. The objective of the unit is to admit care takers, treat/care for the sick, learn, teach practically demonstrations of foods, measurements, preparations and improve the community’s nutrition status of others they live and serve as examples after discharge.

Men put in your time to observe the feeding partens of your children, participate, feed the child, ask what the mother has given prepared and the time of feeds. Help her to prepare the child’s meals. The marriages difference should not interfere with the child’s feeding. Men make a difference in the area of nutrition. Help and strengthen MDG 1, 3, 4, and 5. Mothers alone may find it difficult to take a decision on exclusive breastfeeding without the support of their husbands. With men as members of the support group, it is easier to convince their fellow men in the community as well as to support their wives.

University of Nicaragua

In the country’s “machismo” culture, men control household resources and are not expected to be involved in seeking care for their wives and children, especially during pregnancy, childbirth, and the postpartum period. Women often do not make decisions on their own, which limits their ability to access household financial

Q6 .Do you get involved and participate in feeding your child below 5years at home?

Study area	2003	2007	2016
Katanga peri urban	30%		
Mulago hospital		25%	
Komamboga H/C			50%

The table above showed number of men who said No in the three study populations but accepted to buy foods of their choice, gives some money but not sure of the foods bought and enjoyed by the baby.

MENs INVOLVEMENT AND PARTICIPATION IN FEEDING THE UNDER FIVES;

CAN IT SAVE THE CHILDRENS NUTRITIONALSTATUS ?

Kaggwa Margaret

Uganda

The silent component neglected, " men involvement and participation", untapped targeting only women of child bearing age yet men the policy maker, researchers, bread earner ,administrator, financier, leader ,and God given gift and control all the environment. Malnutrition if not, prevented, treated in early stages promptly, and rehabilitee becomes permanent cause brain damage in the under-fives. Men involved in child neglect, abandonment, divorce, inequalities are leading secondly malnutrition in low income countries. Low adequate feeding practices i.e. continued breast feeding, adequate feeding frequency and diversity in children 6-23 months: 37 percent *ZDHS 2007*). Yet under nutrition is determined by a number of factors which operate at different levels. An individual's nutritional status is influenced by three broad categories of factors -- food, care, and health -- and adequate nutrition requires the presence of all three. Poor infant and young child feeding practices due to lack of resources or knowledge of care takers, along with illnesses such as diarrhea, pneumonia, malaria, and HIV and AIDS, often exacerbated by intestinal parasites, are immediate causes of malnutrition. Underlying and more basic causes include poverty, household food insecurity, unsanitary health environment, illiteracy, social norms, and emergencies In England, the Government's strategy for securing parental involvement was first set out in the 1997 White Paper, Parent & under-fives. Therefore men should be key in reduction to the nutritional status of the under-fives at all levels.

Globally 161 million are under-five have child malnutrition .11 million in Africa and 4 million in Latin America and the Caribbean. High prevalence levels of stunting among children under-five years of age in Africa (United States of America), Ethiopia is one of the countries with the highest levels of malnutrition particularly prevalent among children under five years of age (UNICEF) 2006. The World Health Organization predicts that type 2 diabetes will become the seventh largest killer by 2030. Globally, 1.4 billion people are overweight and obese, while 870 million suffer from chronic hunger, representing a dual burden of malnutrition. Australia is by no means the 'fattest' country. Twenty-five per cent of children in Australia were deemed overweight and obese in 2011-12.

Prevalence of vitamin A deficiency: 54 percent in children under five and 13 percent among women of child bearing age;

Prevalence of anemia: 53 percent in children under five and 30 percent in women of child bearing age (*Source: NFNC2002*)

Under-nutrition can be greatly reduced through the delivery of simple interventions at key stages of the life cycle - for the mother: before she becomes pregnant, during pregnancy, and while breastfeeding; for the child, in infancy and early childhood. Maternal under-nutrition leads to intra-uterine growth retardation and low birth weight. Malnourished children have substantially lower chances of survival than children who are well nourished. They are much more likely to suffer from a serious infection and die from common childhood diseases such as diarrhoea, pneumonia, measles, and malaria. Maternal and child under-nutrition is estimated to contribute to one-third of child mortality.

A package of effective nutrition interventions to reduce the levels of chronic malnutrition (stunting) has been globally agreed upon by experts in the field. It includes: adequate maternal nutrition during pregnancy and lactation, early initiation of breastfeeding, exclusive breast feeding for the first 6 months, continued breastfeeding and adequate complementary feeding from 6 to 24 months, and increased micronutrient intakes during the critical 1,000 days.

UNICEF/World Bank/WHO/GAVI/MoH supports the delivery of priority interventions at different stages of the life cycle: policy development, system strengthening, service delivery, advocacy, capability building, partnerships, resource mobilization, innovations, and knowledge management

Provides technical and financial support to the Ministry of Health under the following programs:
- Bi-annual Child Health Weeks for vitamin A supplementation and de-worming. Children under 5 receive vitamin A capsules and de-worming tablets.

Supports planning, supervision, and implementation.
- Promotion of adequate infant and young child feeding practices through the development of appropriate guidelines and training materials, and capacity building of health workers at facility and community level.
- Management of severe acute malnutrition through the provision of therapeutic nutrition foods, medicines, anthropometric equipment, as well as training of health care workers.

Nutrition is under the program me result, “Children, mothers, and pregnant women benefit from high impact interventions contributing to the attainment of Millennium Development Goal targets for child and maternal survival and development.”

Primary Health Care,Almata Decra lation 1987 targeting by the year 2000 passed unnoticed little was done most communities.

A purposive study was done among men using the same questionnaire in Katanga peri urban slum 2003 (30%), Mulago medical workers 2007 (27%) and at Komamboga rural health center 2016 (50%) about the involvement of males and their participation in feeding the under-fives revealed out that both elite , slum dwellers and rural health workers had never asked the number of times mothers breast feed the baby, all admitted to buy foods in house but didn't know the weaning foods, they admitted to enjoy the fun, love their babies and return home late when babies are asleep rarely earlier than they sleep. At the nutritional Unit of Mulago Kampala national referral hospital out of the 20 admitted cases per day only one mother is being escorted by the husband.

Relapses of malnutrition have been experienced by the unit. The objective of the unit is to admit care takers, treat/care for the sick, learn, teach practically demonstrations of foods, measurements, preparations and improve the community's nutrition status of others they live and serve as examples after discharge.

Men put in your time to observe the feeding partens of your children, participate, feed the child, ask what the mother has given prepared and the time of feeds. Help her to prepare the child's meals. The marriages difference should not interfere with the child's feeding. Men make a difference in the area of nutrition. Help and strengthen MDG 1, 3, 4, and 5. Mothers alone may find it difficult to take a decision on exclusive breastfeeding without the support of their husbands. With men as members of the support group, it is easier to convince their fellow men in the community as well as to support their wives.

University of Nicaragua

In the country's "machismo" culture, men control household resources and are not expected to be involved in seeking care for their wives and children, especially during pregnancy, childbirth, and the postpartum period. Women often do not make decisions on their own, which limits their ability to access household financial

Q6 .Do you get involved and participate in feeding your child below 5years at home?

Study area	2003	2007	2016
Katanga peri urban	30%		
Mulago hospital Komamboga H/C		25%	50%

The table above showed number of men who said No in the three study populations but accepted to buy foods of their choice, gives some money but not sure of the foods bought and enjoyed by the baby.

MENs INVOLVEMENT AND PARTICIPATION IN FEEDING THE UNDER FIVES;

CAN IT SAVE THE CHILDRENS NUTRITIONALSTATUS ?

Kaggwa Margaret

Uganda

The silent component neglected," men involvement and participation", untapped targeting only women of child bearing age yet men the policy maker, researchers, bread earner ,administrator, financer, leader ,and God given gift and control all the environment. Malnutrition if not, prevented, treated in early stages promptly, and rehabilitee becomes permanent cause brain damage in the under-fives. Men involved in child neglect,

abandonment, divorce, inequalities are leading secondly malnutrition in low income countries. Low adequate feeding practices i.e. continued breast feeding, adequate feeding frequency and diversity in children 6-23 months: 37 percent *ZDHS 2007*). Yet under nutrition is determined by a number of factors which operate at different levels. An individual's nutritional status is influenced by three broad categories of factors -- food, care, and health -- and adequate nutrition requires the presence of all three. Poor infant and young child feeding practices due to lack of resources or knowledge of care takers, along with illnesses such as diarrhea, pneumonia, malaria, and HIV and AIDS, often exacerbated by intestinal parasites, are immediate causes of malnutrition. Underlying and more basic causes include poverty, household food insecurity, unsanitary health environment, illiteracy, social norms, and emergencies. In England, the Government's strategy for securing parental involvement was first set out in the 1997 White Paper, *Parent & under-fives*. Therefore men should be key in reduction to the nutritional status of the under-fives at all levels.

Globally 161 million are under-five have child malnutrition .11 million in Africa and 4 million in Latin America and the Caribbean. High prevalence levels of stunting among children under-five years of age in Africa (United States of America), Ethiopia is one of the countries with the highest levels of malnutrition particularly prevalent among children under five years of age (UNICEF) 2006. The World Health Organization predicts that type 2 diabetes will become the seventh largest killer by 2030. Globally, 1.4 billion people are overweight and obese, while 870 million suffer from chronic hunger, representing a dual burden of malnutrition. Australia is by no means the 'fattest' country. Twenty-five per cent of children in Australia were deemed overweight and obese in 2011-12.

Prevalence of vitamin A deficiency: 54 percent in children under five and 13 percent among women of child bearing age;

Prevalence of anemia: 53 percent in children under five and 30 percent in women of child bearing age (*Source: NFNC2002*)

Under-nutrition can be greatly reduced through the delivery of simple interventions at key stages of the life cycle - for the mother: before she becomes pregnant, during pregnancy, and while breastfeeding; for the child, in infancy and early childhood. Maternal under-nutrition leads to intra-uterine growth retardation and low birth weight. Malnourished children have substantially lower chances of survival than children who are well nourished. They are much more likely to suffer from a serious infection and die from common childhood diseases such as diarrhoea, pneumonia, measles, and malaria. Maternal and child under-nutrition is estimated to contribute to one-third of child mortality.

A package of effective nutrition interventions to reduce the levels of chronic malnutrition (stunting) has been globally agreed upon by experts in the field. It includes: adequate maternal nutrition during pregnancy and lactation, early initiation of breastfeeding, exclusive breast feeding for the first 6 months, continued breastfeeding and adequate complementary feeding from 6 to 24 months, and increased micronutrient intakes during the critical 1,000 days.

UNICEF/World Bank/WHO/GAVI/MoH supports the delivery of priority interventions at different stages of the life cycle: policy development, system strengthening, service delivery, advocacy, capability building, partnerships, resource mobilization, innovations, and knowledge management

Provides technical and financial support to the Ministry of Health under the following programs:
- Bi-annual Child Health Weeks for vitamin A supplementation and de-worming. Children under 5 receive vitamin A capsules and de-worming tablets.

Supports planning, supervision, and implementation.
- Promotion of adequate infant and young child feeding practices through the development of appropriate guidelines and training materials, and capacity building of health workers at facility and community level.
- Management of severe acute malnutrition through the provision of therapeutic nutrition foods, medicines, anthropometric equipment, as well as training of health care workers.

Nutrition is under the program me result, “Children, mothers, and pregnant women benefit from high impact interventions contributing to the attainment of Millennium Development Goal targets for child and maternal survival and development.”

Primary Health Care,Almata Decra lation 1987 targeting by the year 2000 passed unnoticed little was done most communities.

A purposive study was done among men using the same questionnaire in Katanga peri urban slum 2003 (30%), Mulago medical workers 2007 (27%) and at Komamboga rural health center 2016 (50%) about the involvement of males and their participation in feeding the under-fives revealed out that both elite , slum dwellers and rural health workers had never asked the number of times mothers breast feed the baby, all admitted to buy foods in house but didn’t know the weaning foods, they admitted to enjoy the fun, love their babies and return home late when babies are asleep rarely earlier than they sleep. At the nutritional Unit of Mulago Kampala national referral hospital out of the 20 admitted cases per day only one mother is being escorted by the husband. Relapses of malnutrition have been experienced by the unit. The objective of the unit is to admit care takers, treat/care for the sick, learn, teach practically demonstrations of foods, measurements, preparations and improve the community’s nutrition status of others they live and serve as examples after discharge.

Men put in your time to observe the feeding partens of your children, participate, feed the child, ask what the mother has given prepared and the time of feeds. Help her to prepare the child’s meals. The marriages difference should not interfere with the child’s feeding. Men make a difference in the area of nutrition. Help and strengthen MDG 1, 3, 4, and 5. Mothers alone may find it difficult to take a decision on exclusive breastfeeding without the support of their husbands. With men as members of the support group, it is easier to convince their fellow men in the community as well as to support their wives.

University of Nicaragua

In the country’s “machismo” culture, men control household resources and are not expected to be involved in seeking care for their wives and children, especially during pregnancy, childbirth, and the postpartum period. Women often do not make decisions on their own, which limits their ability to access household financial

Q6 .Do you get involved and participate in feeding your child below 5years at home?

Study area	2003	2007	2016
Katanga peri urban	30%		
Mulago hospital		25%	
Komamboga H/C			50%

The table above showed number of men who said No in the three study populations but accepted to buy foods of their choice, gives some money but not sure of the foods bought and enjoyed by the baby.

key in reduction to the nutritional status of the under-fives at all levels.

Globally 161 million under-five are malnourished. High incidence in Africa 11 million and 4 million in Latin America and the Caribbean. Ethiopia is one of the countries with the highest levels of malnutrition particularly prevalent among children under five years of age (UNICEF) 2006. The World Health Organization predicts that type 2 diabetes will become the seventh largest killer by 2030. Globally, 1.4 billion people are overweight and obese, while 870 million suffer from chronic hunger, representing a dual burden of malnutrition. Australia is by no means the 'fattest' country. Twenty-five per cent of children in Australia were deemed overweight and obese in 2011-12.

Prevalence of vitamin A deficiency: 54 percent in children under five and 13 percent among women of child bearing age;

Prevalence of anemia: 53 percent in children under five and 30 percent in women of child bearing age (*Source: NFNC2002*)

Under-nutrition can be greatly reduced through the delivery of simple interventions at key stages of the life cycle - for the mother: before she becomes pregnant, during pregnancy, and while breastfeeding; for the child, in infancy and early childhood. Maternal under-nutrition leads to intra-uterine growth retardation and low birth weight. Malnourished children have substantially lower chances of survival than children who are well nourished. They are much more likely to suffer from a serious infection and die from common childhood diseases such as diarrhea, pneumonia, measles, and malaria. Maternal and child under-nutrition is estimated to contribute to one-third of child mortality.

A package of effective nutrition interventions to reduce the levels of chronic malnutrition (stunting) has been globally agreed upon by experts in the field. It includes: adequate maternal nutrition during pregnancy and lactation, early initiation of breastfeeding, exclusive breast feeding for the first 6 months, continued breastfeeding and adequate complementary feeding from 6 to 24 months, and increased micronutrient intakes during the critical 1,000 days.

UNICEF/World Bank/WHO/GAVI/MoH supports the delivery of priority interventions at different stages of the life cycle: policy development, system strengthening, service delivery, advocacy, capability building, partnerships, resource mobilization, innovations, and knowledge management

Provides technical and financial support to the Ministry of Health under the following programs:
- Bi-annual Child Health Weeks for vitamin A supplementation and de-worming. Children under 5 receive vitamin A capsules and de-worming tablets.

Supports planning, supervision, and implementation.
- Promotion of adequate infant and young child feeding practices through the development of appropriate guidelines and training materials, and capacity building of health workers at facility and community level.
- Management of severe acute malnutrition through the provision of therapeutic nutrition foods, medicines, anthropometric equipment, as well as training of health care workers.

Nutrition is under the program me result, “Children, mothers, and pregnant women benefit from high impact interventions contributing to the attainment of Millennium Development Goal targets for child and maternal survival and development.”

Primary Health Care, Almata Decra lation 1987 targeting by the year 2000 passed unnoticed little was done most communities.

A purposive study was done among fathers using the same questionnaire in Katanga peri urban slum 2003 (30%), Mulago medical workers 2007 (27%) and at Komamboga rural health center 2016 (50%) about the involvement of males and their participation in feeding the under-fives revealed out that both elite , slum dwellers and rural health workers had never asked the number of times mothers breast feed the baby, all admitted to buy foods in house but didn’t know the weaning foods, they admitted to enjoy the fun, love their babies and return home late when babies are asleep rarely earlier than they sleep. At the nutritional Unit of Mulago Kampala national referral hospital out of the 20 admitted cases per day only one mother is being escorted by the husband. Relapses of malnutrition have been experienced by the unit. The objective of the unit is to admit care takers, treat/care for the sick, learn, teach practically demonstrations of foods, measurements, preparations and improve the community’s nutrition status of others they live and serve as examples after discharge.

Fathers put in your time to observe the feeding par tens of your children, participate, feed the child, ask what the mother has given prepared and the time of feeds. Help her to prepare the child’s meals. The marriages difference should not interfere with the child’s feeding. Men make a difference in the area of nutrition. Help and strengthen MDG 1, 3, 4, and 5. Mothers alone may find it difficult to take a decision on exclusive breastfeeding without the support of their husbands. With men as members of the support group, it is easier to convince their fellow men in the community as well as to support their wives.

University of Nicaragua

In the country’s “machismo” culture, men control household resources and are not expected to be involved in seeking care for their wives and children, especially during pregnancy, childbirth, and the postpartum period. Women often do not make decisions on their own, which limits their ability to access household financial

Q6 .Do you get involved and participate in feeding your child below 5years at home?

Study area	2003	2007	2016
Katanga peri urban	30%		
Mulago hospital		25%	
Komamboga H/C			50%

The table above showed number of fathers who said No in the three study populations but accepted to buy foods of their choice, gives some money but not sure of the foods bought and enjoyed by the baby.